

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 47/10, 47/38, 9/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/25544</b> <b>(43) International Publication Date:</b> 28 September 1995 (28.09.95)
<b>(21) International Application Number:</b> PCT/EP95/01025 <b>(22) International Filing Date:</b> 20 March 1995 (20.03.95) <b>(30) Priority Data:</b> 0325/94                      21 March 1994 (21.03.94)                      DK <b>(71)(72) Applicant and Inventor:</b> BROWN THOMSEN, John [FR/FR]; "La Campagne", 587, chemin du Clot, F-06510 Gattières (FR). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> MØLLER, Jens, Christian [DK/DK]; Osterbrogade 24, DK-7620 Lemvig (DK). <b>(74) Agent:</b> GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
<b>(54) Title:</b> GEL FOR TREATMENT OF SKIN DISEASES AND FOR DISINFECTION OF THE SKIN  <b>(57) Abstract</b>  Gel for local treatment of skin diseases and for prophylaxis, characterised by containing more than 90 % of a drying and/or protein coagulating, short-chained alcohol or alcohol mixture, primarily ethanol, and possibly adjuvants or additives and by containing a gelling agent, that possesses good skin-adhesive properties, that gives a matrix formation of alcohol or alcohol mixtures, that creates an evaporation inhibiting effect, gives a prolonged effect, and form a protective plaster when the gel has dried.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

GEL FOR TREATMENT OF SKIN DISEASES  
AND FOR DISINFECTION OF THE SKIN

This application concerns a matrix-forming and  
5 skin-adhesive, anti-evaporant gel for local treatment of  
skin diseases, for both primary and secondary prophylactic  
treatment of wounds, and for skin disinfection or the like  
in both humans and animals. More precisely, the invention  
relates to a gel characterized by containing lower alkanol  
10 in a concentration of more than 90% and by containing a  
gelling agent, such as ethyl(hydroxyethyl)cellulose or  
another suitable gelling agent, and possible additives,  
whereby the gel can be applied to the skin efficiently,  
easily, and without complications.

15

BACKGROUND

Infectious skin diseases, both microbial and  
parasitic, are widespread. Some examples of skin diseases  
are viral skin infections caused by, for example, Herpes  
20 simplex virus or Varicellae Zoster virus, bacterial skin  
infections caused by, for example, Staphylococcus aureus,  
fungal infections caused by, for example, Trichophyton  
rubrum, for eliminating skin parasites, such as Sarcoptes  
scabiei var. hominis. The virus infection, Herpes simplex,  
25 alone occurs with approximately 100 million new cases per  
year.

Herpes simplex is caused by Herpes simplex virus  
(HSV). This virus occurs in two antigenically different  
types, type 1 and type 2. Herpes on the lips and around  
30 the mouth (Herpes labialis) is usually caused by type 1;  
most incidences of Herpes on and around the genitals  
(Herpes genitalis) are caused by type 2.

The first infection with HSV (primary infection)  
varies symptomatologically. Usually it occurs during  
35 childhood. At this first infection, HSV-DNA is  
incorporated into the cells. Intermittently, virus  
proliferation occurs, resulting in Herpes outbreaks from

this incorporated HSV-DNA (these outbreaks are called secondary infections). Most Herpes outbreaks in adults are secondary, where the infection flares up due to reduced resistance, febrile diseases (e.g. Pneumonia), traumas, or  
5 the effects of cold, heat, or light.

The outbreak starts with flushing, swelling, itching, and pain in the infected area followed by necrosis and a suppurative ulceration which is the most troublesome symptom. A spontaneous healing of the wounds will occur in  
10 approximately 10 to 14 days.

The outbreak of Herpes genitalis - mentioned above - is identical to that of Herpes labialis except for the fact that Herpes genitalis is located on and around the genitals.

15 The number of therapeutics for, for example, HSV skin infections is very limited, and the present antiviral chemotherapy has not convincingly been proved efficient. Furthermore, there exists a few products for advancing the healing of established HSV-wounds and inhibition of further  
20 outbreaks. However, none of these products have a convincing effect.

Bacterial infections such as pimples, isolated or in connection with Acne vulgaris, are examples of other skin diseases for which adequate treatments are not currently  
25 available. In Acne it is currently believed that lipolysis of triglycerides by Propionibacterium acnes releases fatty acids; it is thought that these fatty acids are capable of eliciting an inflammatory process in the follicle wall, which may then rupture. Discharge of the follicular  
30 content leads to perifollicular inflammation. Benzoyl peroxide, which is currently used for treating acne, has a mode of action dependent upon its broad-spectrum antimicrobial effect on both primary inflammation and on secondary infection. Current treatments based on benzoyl  
35 peroxide have inherent problems, including lack of efficacy and adverse reactions such as erythema, burning and desquamation of the skin.

From the literature it is known to use alcohol as disinfectant against, for example, virus including HSV - see, for example, R. Tyler; Journal of Hospital Infection (8: 22-29; 1987). When using alcohols as normal liquids a  
5 poor and very brief effect is achieved due to the very rapid evaporation of the alcohols. Also the use of alcohols at concentrations below 90% by weight gives inadequate results.

Furthermore, Moldenhauer, in Zbl. Bakr. Hyg., I Abt.  
10 Orig. B 179, 544-554 (1984) compares surface disinfection properties of ethanol, isopropanol, formaldehyde and benzalkonium chloride by suspending virus suspension (including HSV, influenza, cocksackie-B and mumps) in those compounds or solutions. Alcohol concentrations above 90%  
15 were not tested. Furthermore in these two references alcohol is being used for surface disinfection properties and not for treatment of infections and the symptoms thereof.

In US 5,145,663, a disinfectant, consisting of 65-75%  
20 isopropyl alcohol, 8-12% propylene glycol, and potential inert ingredients or disinfectants or antiseptics, is mentioned. The patent does not mention gels.

In GB-A-2017491 a gel containing alcohol is used as a hand-wash for bacterial disinfection.

25 In the above references describing the use of alcohol, either as such or as a solvent for other active disinfectant agents, in surface disinfection of the skin, the alcohol will remain in contact with the skin for a relatively short period of time. The compositions, if they  
30 are washing compositions when they may contain a thickener such as sodium chloride, are generally rinsed off with water. Treatment with no-rinse compositions and alcohol wipes applies a relatively low amount of composition and the alcohol evaporates quickly. The alcohol is, in  
35 particular, not in contact with the skin long enough for penetration to layers below the stratum corneum (dermis and epidermis).

Ethanol has been used extensively in pharmaceutical compositions used for topical application to the skin. Compositions including a gelling agent have, likewise, been used for such application. However in none of these disclosures is the alcohol itself used as an agent for treating skin diseases. The following references are relevant.

In US 3,016,328 and in US 4,590,214 a mixture of a dialdehyde and an alcohol is mentioned. Without evaluating the effect, it can be established that none of these products include alcohol and gel-forming agents.

WO 93/00114 describes a method for reducing the duration of HSV-infection by applying a mixture of an anaesthetic and a surface-active ingredient with suitable antiviral activity. It does not mention gelling agents.

US 4,247,547 mentions the use of gels containing alcohols and the dermatologically active tretinoin for treatment of acne. Tretinoin is an irritant and the compositions would be wholly unsuitable for treatment of skin infected by HSV. Also the concentration of water in the compositions is unclear.

In Chemical Abstracts 90:76564r (1979) an antiseptic paste is disclosed containing about 80% by weight ethanol, 13% water and a thickener.

US 5,013,545 describes a gel consisting of 60-90% ethanol, 0.5-30% water, and an active ingredient, such as an antihistamine. This patent does not mention activity on viral skin infections such as Herpes. Further, none of the worked examples teaches how to produce a stable gel with more than 80% alcohol.

US 5,098,717 describes a gel based on 60-90% ethanol and as active ingredients an antihistamine and an antipruritic.

Carrier gels for pharmaceuticals based on ethanol and water are described in the patent literature, see, for example, SE 466134. In US 4,593,048 it is mentioned that the penetration into the circulation by pharmaceuticals,

dissolved in ethanol and applied topically, is accelerated when various adjuvants are used. The formulation contains surface-active ingredients as penetration aids for pharmaceuticals for percutaneous systematic administration.

5 The compositions are not used to treat skin disorders.

An article by B. Rodu and F. Lakeman ("In vitro virucidal activity by component of a topical film-forming medication," J.Oral Pathology 17: 324326; 1988) mentions in vitro trials of a preparation consisting of approximately  
10 80% ethanol, 7% tannic acid, 2.5% salicylic acid, and 1% boron acid. The tests were intended to evaluate the in vitro properties of the product Zilactin, which contains those ingredients and a hydroxy propylcellulose gelling agent. The in vivo performance of the gel against HSV has,  
15 however been found to be limited.

A specific antiviral preparation for topical treatment of Herpes is Zovir/Zovirax cream (Zovir is a registered trademark), which contains 5% aciclovir. The efficacy has not been convincingly proved in relation to Herpes simplex  
20 skin infections, and, moreover, reports that resistance is beginning to develop have been published.

Acyclovir has been administered systemically for the treatment of Varicellae Zoster virus. However reports have indicated that the treatment limited the immune response  
25 with consequent failure to develop resistance to the disease.

Surprisingly, it has now turned out that it is possible to produce an effective gel for treating skin diseases and for controlling skin parasites without using  
30 anti-histamines, anaesthetics, anti-inflammatory agents, and totally without using pharmaceuticals, including biocides against skin parasites.

Ethyl hydroxy ethylcellulose (EHEC) is produced by a first swelling native cellulose in alkali, then adding  
35 ethylene oxide to cellulose hydroxyl groups activated in the first step, then etherifying hydroxyl groups in the product by reacting them with ethyl chloride after alkali

treatment. In the ethylene oxide treatment step ethoxy units may be added to the hydroxyl group on a pendant group derived from the earlier reaction of an ethylene oxide molecule with a cellulose-hydroxyl group. In the etherification step, hydroxyl groups of pendant groups and of cellulose-hydroxyl group may be reacted. The polymer product thus contains ethoxy 2-ethoxyethyleneoxy and ethoxypoly(ethyleneoxy) pendant groups. The reactions can be controlled so as to provide EHEC products with a variety of different degrees of substitution, and molar substitution (i.e. a measure of the average ethyleneoxy units per etherified group). These parameters, as well as the degree of polymerisation/ molecular weight, affect the properties of the polymer in solution.

The performance of EHEC in aqueous systems, where it is used as a thickener and dispersing agent, for instance in paints and cement based mortar, has been studied (Törnquist, J, Farg och Lack Scandinavia, 31, 291-295 (1985), Carlsson, A et al polymer, 27, 431-436 (1986)). We are not aware that its use to thicken or gel non-aqueous alcohols has been studied.

#### SUMMARY OF THE INVENTION

A new gel-form pharmaceutical composition according to the invention comprises a liquid and a high molecular weight polymer gelling agent dissolved in the liquid, characterised in that the composition comprises more than 90% by weight of a lower alkanol based on the total weight of the composition and less than 10% by weight water based on the total weight of the composition.

The composition optionally comprises one (or more) enhancing agent which enhances the effect of the alcohol. Preferably the composition is substantially free of antihistamines, anaesthetics, anti-inflammatories, irritants and any immunogenic compounds or compounds which disturb the immune system. Consequently, in the invention, concentration and crystallization of medicaments (which is a potential problem with prior art compositions in which



alcohol is held as a solvent vehicle for such active compounds) will not occur when the solvent evaporates. Consequently, local overdosing resulting in irritative inconveniences is totally avoided.

5       The lower alkanol preferably contains up to 4, preferably up to 3 carbon atoms. It may be a glycol or polyol but is preferably a mono hydroxy compound. It can be a mixture of such compounds. Most preferably it includes ethanol optionally in combination with other lower  
10       alkanols.

Thus, it has now surprisingly been found that a gel containing more than 90% ethanol or other lower alkanol is very effective for local treatment of, for example, skin infections and skin parasites.

15       When using suitable gelling agents it is possible to transform a concentrated alcohol into a very suitable, effective, and stable gel. The invention is of particular value in the treatment of viral infections whose systems involve skin eruptions, especially herpes infections.

#### 20       DETAILED DESCRIPTION OF THE INVENTION

Thus, the preparation of the invention consists of concentrated alcohol in such a form that it is usable as a topical preparation for immediate application to the infected area of the skin including mucous membranes.  
25       While ethanol in lower concentrations, for example in concentrations of less than 60%, gives a distinct pain reaction, the use of the concentrated alcohol, for example in concentrations of more than 90% according to the invention, is almost painless even when it is used in open  
30       wounds. The preparation has a combined effect which is utilised optimally and over a prolonged period in the invention. Thus, an effective combination of drying the edematous tissue, coagulating proteins and killing the infectious agent, is achieved, and later, when it has  
35       dried, the gelling agent acts as a plaster protecting against infection.

The gels of the invention create a matrix formation that prevents the alcohol from leaking out of the gel and flowing away from the site where the gel has been applied. At the same time the gel is gentle to the skin and easy to  
5 apply. It is possible, and preferred, for the composition to be transparent, and for the dried film of polymer gelling agent, remaining after the composition has dried on the skin, to be transparent to allow the underlying skin to be visible throughout treatment.

10 When the gel has dried it forms a protective film over a wound such as an HSV outbreak. Separately, all these activities contribute to avoid secondary bacterial infection. With the evaporation-limiting matrix formation the gel forms a barrier film on the surface between gel and  
15 air by means of which the evaporation of alcohol is drastically reduced. For a long time the alcohol in this way remains in contact with the skin by which means the ethanol has a possibility of diffusing into the skin and perform its effect in depth. Further, the plaster action  
20 of the gel prevents a rapid evaporation of the alcohol diffused into the tissue and, finally, after drying the plaster effect of the gel will offer protection against reinfection of the affected area. Thus, when choosing a gelling agent or a mixture of gelling agents, which form  
25 matrix with the ethanol and that form a film on the surface of the gel against the atmosphere, the extraordinary prolonged effect of the gel is achieved according to the invention..

The content of water is important to the  
30 effectiveness. The content should be less than 10%, preferably less than 5%, and optimally, in some cases, less than 1%. The amount of water should be below the equilibrium content of the composition under ambient conditions, that is under normal storage conditions at 20-  
35 24°C and at 50-100% relative humidity, as well as at such humidity levels at temperatures up to body temperature (eg 37°C). Thus the composition as a whole should, in effect

be hygroscopic. The hygroscopicity is preferably due to the concentration of alkanol being higher than the concentration of alkanol in the presence of humid air. The equilibrium concentration of water in ethanol under these conditions is in the range 3 to 7% by weight of the sum of water and ethanol. The gelling agent may also contribute some hygroscopicity. If the content of water is too high the drying and antimicrobial effect and especially the antiviral effect is reduced which means that the effectiveness of the gel is reduced. Similarly, the concentration of alkanol is crucial to the effect on external skin parasites. It seems that the high concentration of alkanol may allow the alkanol to penetrate to layers of skin below the stratum corneum.

One of or a combination of several gelling agents which are usually soluble in the alkanol can be used. The gelling agent is a polymer, which may be linear, branched or cross-linked and may be naturally-derived, a derivative of such polymers, or may be wholly synthetic. The molecular weight is usually high, for instance at least  $10^4$ , preferably at least  $5 \times 10^5$ , and up to several million, for instance more than  $10^6$ . The polymer is preferably non-ionic in the composition. Suitable gelling agents include cellulose derivatives, especially cellulose ethers, such as ethylcellulose, methylcellulose; hydroxyethyl cellulose especially ethyl(hydroxyethyl)cellulose (EHEC), alkyl- and hydroxyalkyl cellulose, carboxy methyl cellulose, other polysaccharides, such as naturally derived polysaccharides and derivatives thereof, including modified carragenan; and synthetic polymers such as polyethylene glycols, polyethylene oxides, polyvinyl pyrrolidones and polyacrylic acid.

The cellulose derivatives which are useful in the invention usually have a high molecular weight, for instance more than  $10^6$ , although polymers with molecular weight from  $10^5$  upwards may be useful. The degree of substitution/derivatisation of cellulose ether derivatives

useful in the invention is preferably relatively high, for instance higher than 1.0.

A particularly suitable cellulosic gelling agent for a highly concentrated ethanol is EHEC, ethyl(hydroxyethyl)cellulose, which is a derivative of cellulose with CA registration number 9004-58-4. EHEC is, as an example, sold under the trademark BERMOCOLL from Berol Kemi AB, such as BERMOCOLL OS. For example, an effectively gelling EHEC is achieved at degree of polymerisation of approximately 3,200, a degree of substitution of approximately 1.7 for ethylene (DS-ethyl = 1.7) and of approximately 1.5 for hydroxyethyl (MS-hydroxyethyl = 1.5). Celluloseethers including EHEC are more precisely described in Kirk-Othmer, "Encyclopedia of Chemical Technology," 5:143,1979(3. edition). See also "Faerg och Lack Scandinavia" 31:291-298;1985.

Acrylic acid polymerics are also particularly suitable as gelling agent. Acrylic acid polymerics are, as an example, sold under the trademark Carbopol from BF Goodrich, such as Carbopol 940 and 941, Carbopol 940 NF and 941 NF, Carbopol 980 NF and 981 NF, or Carbopol 1342 and 1382. Those Carbopols are high molecular, non-linear polymerics of acrylic acid cross-linked with polyalkenyl polyether. Acrylic acid polymerics are more precisely described in "Kirk-Othmer, Encyclopedia of Chem.Tech," 20:216;1982 and in Ullmanns Encyclopedia of Ind.Chem," A21:752;1992.

Polyvinylpyrrolidones are a third example of a particularly suitable gelling agent. Polyvinylpyrrolidones, as an example, are sold under the trademark PVP K-30 and PVP K-90 from GAF. Polyvinylpyrrolidones are high molecular polymerics which are described in more detail in "Kirk-Otmer, Encyclopedia of Chem.Tech.," 23:963;1983 and in "Ullmanns Encyclopedia of Ind.Chem.," A21:143;1992 and others.

The gelling agents are used in amounts between 0.1% and 10%, depending on the choice of gelling agent or

mixture of gelling agents, depending on the composition, the desired texture etc. The amount should preferably be sufficient to render the composition gel-like at room temperature and at normal body temperature so that it remains in place on the skin and does not spread or run off after application. The gel-like consistency, which is due to the viscoelastic properties of polymer solutions in solvents, depends upon molecular weight, degree of substitution as well as, for derivatives such as EHEC where the several units of derivatising agent may be added to each derivatised saccharide hydroxyl group, the molar substitution and type of substituent as well as the concentrations of polymer in the composition. The compositions should generally have high viscosity under low shear but, for optimal handling during manufacture and application, should be shear thinning. This combination of features can be achieved by appropriate selection of properties, as illustrated in the accompanying examples, for instance. For example, the viscosity-increasing effect of EHEC depends on the degree of polymerization and on the degree of substitution and, for the EHEC having a degree of polymerisation of 3200, mentioned above, a suitable concentration in ethanol is 0.5 to 2.0%, for instance around 1.0%.

This, for example, applies to pH-regulating agents such as bases, eg alkaline inorganic compounds or organic bases and mixtures by which the action of the alcohol is increased under certain circumstances. Inorganic bases which may be used include sodium and potassium hydroxide and carbamate. Organic bases include triethylamine, triethanol amine and other alkanolamines. For example, a content of 0.02% NaOH will increase the antiviral action of the ethanol. Thus in one embodiment of the invention the composition has a pH in the range 6 to 9.5, preferably an alkaline pH. Other additives that can be mentioned are the substances that form part of medicinal gels such as emollients, colorants, perfumes, menthol, camphor,

W-protective agents etc. and the like by which the gel can be supplemented with further functional properties.

The composition should, however, be substantially free of pharmacologically active ingredients other than these optional enhancing agents.

The composition is preferably supplied in an air and moisture/moisture vapour-impermeable container. Such containers are, for instance, squeezable tubes, especially formed of metal foils or of plastics materials having moisture barrier properties. Such containers prevent compositions, whose water content is such that the composition is hygroscopic, from absorbing moisture from the atmosphere during storage and before use. By the use of such containers, therefore loss of drying activity of the gels is minimised.

According to the invention we have not only succeeded in producing a gel - with a concentrated content of alkanol - which is very effective and suitable for treatment of skin diseases, which is skin-adhesive and gentle to the skin, and which preferably does not contain other medicaments or pharmaceuticals. The mere omission of medicaments and pharmaceuticals such as antihistamines etc. has as a consequence that no reverse actions or side-effects occur and that allergic reactions are completely eliminated. Further to this, as a consequence of the special mechanism of activity of ethanol, absolutely no resistance can develop among the responsible microorganisms or parasites. In choosing a gel with a matrix structure it is achieved that the ethanol, after application, does not accumulate in, for example, the nasolabial fissure at the angle of the mouth or in the groin but remains where applied. Further, as the gel shows pseudoplastic (viscoelastic) properties the gel is very easy to apply, and at the same time it regains its matrix structure and its structural firmness and exactly by that, as mentioned, remains on the site of application.

Because of the high concentration of alcohol the gel possesses other surprising properties. The skin-adhesive properties turn out to be very good, partially due to, the high content of alkanol, but also because of the choice of gelling agent, where especially EHEC, ASP, and PVP or combinations of these have lipophilic and hydrophilic properties giving the alcohol a very good contact with the skin.

The gelling agents, especially EHEC, ASP, and PVP or combinations of these, have hydrophobic-hydrophilic properties by which the release of alcohol towards the skin from the slow release matrix structure of the gel is adjustable. By doing this, it can be obtained that no release-inhibiting film is created between the alcohol-gel and the skin/mucous membranes. This, of course, is important for the continuous effect of the alcohol-gel on the site of application.

Thus, it is not necessary to add surfactants to achieve the correct contact between the skin and the gel as is the case in US 4,593,048. The composition should generally be free of added surfactants.

Similarly, it is not necessary to add special binding agents nor to use plaster or tape in order to adhere the gel to the skin.

Gels according to the invention are physically and chemically stable for at least 12 months at 50°C. Among other things this is a result of the fact that addition of other active ingredients or adjuvants, which together with medicament or biocide may be labile during production and storage, is not needed. It is not necessary to add actual medicaments that, in turn, would require protective antioxidants etc. to secure the chemical stability of the very same medicaments during production and storage.

As the gel does not contain actual skin irritants it is, as mentioned, not necessary to add anti-irritants such as anti-histamines, anti-inflammatory agents or similar agents.

Because of the concentrated content of alcohol the gel is self-preserving. It is therefore not necessary to add antimicrobial preservatives against fungoid growths nor products against bacteria or other micro-organisms, and it is not necessary to store the gel in refrigerator or the like. Omitting all these additives in the gel means that undesirable side-effects of such additives are eliminated.

Two very important properties are achieved in the invention by omitting surfactants, skin adhesives, pharmaceuticals, medicaments, antioxidants, antihistamines, or other anti-inflammatory agents, and because it is not necessary to add preservatives against fungi, bacteria, or other micro-organisms to the gel. First, in all simplicity, the gel is composed of non-allergenic substances. Secondly, owing to its special mode of action towards the infectious agents ethanol does not give rise to development of resistance. Furthermore the product does not disturb the hosts immune response.

For the sake of completeness it should be mentioned that many skin diseases, for example, Herpes simplex are complicated by secondary, usually bacterial infections. It is not necessary to add other pharmaceuticals to the gel to avoid secondary infections, as the alcohol in the gel with the long-term effect disinfects the area and protects it against reinfection via the plaster effect of the dried gel until the wound has healed.

Especially regarding Herpes infections the drying and protein coagulating effect is of great importance. During the primary phase of the repeated herpes outbreaks (secondary infections) are characterized by the formation of blisters, full of liquid. Apart from an immediate improvement of the itching and the pain reaction, a drying action on the blisters, that have already formed, is achieved when the preparation is applied at an early stage, and thus the blisters will disappear rapidly. If the herpes outbreak is not treated early enough the blisters will burst resulting in the formation of suppurating, open



wounds that are characteristic of the secondary phase. When used during this phase, the preparation has an immediate drying effect by means of which the secretion ends, and the protein coagulating action destroy the superficial, necrotic cells. The drying effect makes the product useful on moist skin areas. Here the drying effect has an immediate prophylactic action in relation to bacterial infections through a reduction of the growth conditions.

Similar advantages are seen when the gel is used to treat skin eruptions associated with other viral infections such as chickenpox. When applied to such skin eruptions the formation of blisters is prevented or they are dried and itching is minimised. The breaking of the skin with subsequent risk of infection is thus prevented.

In this connection we can also mention accidental skin injuries such as wounds and skin abrasions where the antimicrobial effect combined with the plaster action of the gel after drying has an indisputable preventive action in relation to wound infections.

The gel is further suitable for prophylactic treatment of physical skin injuries such as cuts, abrasions etc. This indication is not only based on the antimicrobial property of the preparation, but also on the protective effect of the dried gel having the "built-in" plaster effect that is achieved when the preparation has dried.

It is believed that use of the preparation in connection with treatment of burns will have a useful infectious prophylactic effect. Some eczematous diseases including allergic skin diseases are characterized by secondary infections. Here the invention will have great importance too, not least because of the non-allergenic property of the composition.

As mentioned above the gel is suitable for controlling externally parasitical and troublesome organisms. For example, the gel is suitable for external treatment of scabies, chigger and other ectoparasites.

A very effective preparation against skin infections and for eliminating external skin parasites can be obtained when using a gel in which the liquid consists of concentrated ethanol or a concentrated mixture of ethanol and other short-chained alcohols such as isopropanol or propylene glycol and containing in addition, additives such as drying and disinfecting agents, as well as the gelling agent, but no other ingredients.

The gel is also found to be useful for application to the site of insect bites and stings. The effect is thought to be due at least partially to the function of the concentrated alcohol in causing toxic proteins in the sting to coagulate and be rendered inactive.

The invention is further illustrated in the following examples.

#### EXAMPLES

Examples of compositions of the gel:

##### Example 1

	<u>Ingredient</u>	<u>Weight in %</u>
20	Ethanol 99.9%	91
	Water	7
	Carbopol 980 NF	2

Carbopol 980 NF (trade mark, BF Goodrich) has a molecular weight of around  $7.5 \times 10^5$ .

##### 25 Example 2

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol 99.9%	95
	Water	3
	Carbopol 980 NF	2
30	Perfume	-

##### Example 3

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol 99.9%	min. 99
35	Carbopol 940 NF*	max.

\* Carbopol 940 NF (registered trademark) an acrylic acid polymer from BF Goodrich

Example 4

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol 99.9%	97.3
	Bermocoll OS*	2.3
5	NaOH or triethylamine	0.1
	Water	0.3

\* Bermocoll OS (registered trademark) from Berol-Nobel  
= Ethyl(hydroxyethyl)cellulose

10 Example 5

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol 99.9%	91.0
	EHEC*	1.5
	Glycerine	1.0
15	Propyleneglycol	3.0
	Water	3.5

\*(DP = 1,600, DS-ethyl = 0.8, MS-hydroxyethyl = 2.0)

Example 6

	<u>Ingredient</u>	<u>Weight in %</u>
20	Ethanol 99.9%	91.0
	PVP K-30*	2.5
	UV-absorbent	1.0
	Camphor	0-0.5
25	Propylene glycol or adjuvant	0-3.0
	Water	0-2.0
	Polyvinyl alcohol	0-1.0

\* PVP K-30 (registered trademark) from GAF = Polyvinylpyrrolidone, average molecular weight of  $4 \times 10^4$ .

30

Example 7

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol	91
	Carbopol 940	1
35	PVP K-90	2
	Water	4
	Surfactant	2

Example 8

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol	91.0
	Hydroxypropylcellulose	2.5
5	Xanthan gum	0.1
	Isopropanol	4.0
	Water	2.4

The hydroxypropyl cellulose is selected for its solubility in the liquid.

10 Example 9

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol	93.0
	Hydroxyethylcellulose	1.0
	Ethylcellulose	0.5
15	Methylcellulose	0.5
	NaOH	0.02
	Water	4.98

All the cellulose derivatives are selected for their solubility in the liquid.

20 Example 10

	<u>Ingredient</u>	<u>Weight in %</u>
	Ethanol	95.0
	Ethylhydroxyethylcellulose (Bermocoll OS)	1.0
	Polyvinylpyrrolidone (MW $3.5 \times 10^4$ )	2.2
25	Polyethyleneglycol (MW $2 \times 10^3$ )	0.2
	KOH	0.02
	Water	1.58

Examples of the use of the gels:

30 Example 11Local treatment of Herpes simplex

When used for local treatment of Herpes simplex the effect depends on the time of application in relation to the start of the infection. Gels with particularly high concentration of alcohol are suitable, for example, gels mentioned in example 3, 4 or 10.

Within the first hours of the first phase which is characterized by itching, slight pain and prickling, followed by formation of vesicles, the gel is applied to the affected area approximately once an hour. The  
5 subjective symptoms dwindle immediately, and after 3-5 applications the outbreak is stopped and the vesicles disappear. After this, the gel is re-applied every 3-5 hours for 24 hours to secure that the outbreak has stopped completely.

10 When treatment is not started before the vesicles have burst and the gel is applied with a frequency of about every 3 hours the wounds are kept dry without the suppuration. The treatment is continued with decreasing application frequency until all wounds have started  
15 healing.

All three gels gave good results when used at these stages of the infection.

#### Example 12

##### Treatment of Herpes zoster

20 When used for treating local skin outbreaks of Herpes zoster the gel mentioned in example 2 and 9 have been applied to the affected area. During the first 24 hours the gel is applied approximately every 2 hours, and after that at longer intervals as required. The vesicles heal up  
25 and the patient's inconveniences will abate rapidly.

#### Example 13

##### Treatment of bacterial skin infections

The treatment of skin infections in connection with Acne vulgaris should be mentioned as an example of treating  
30 bacterial skin infections.

Gel in accordance with examples 1 and 6 according to the invention is applied to the infected skin areas. To begin with the gel is applied 3 times a day, and after that, when the infected areas have dried it is applied  
35 approximately once or twice a day. By means of this treatment with each of the gels rapid healing is achieved.

A gel, in accordance with example 3, can be used in difficult cases directly on extensively infected areas.

Example 14

5 Treatment of mycologic skin infections

The treatment of epidermophytosis (Tinea pedis) should be mentioned as an example of treating mycologically conditioned infections. Rapid healing of fissures, and successive cure of the fungus infection is achieved when a  
10 gel, in accordance with example 4, 7 and 8, is applied twice a day primarily, and after a few days only once a day.

Example 15

15 Control of external parasites

As required a thin or thick layer of a gel, in accordance with example 3 and 9 of the invention, is applied to the infected area with the external parasites such as lice, scab mites, ticks and crab lice. In a short  
20 time the parasites are eliminated.

Because of the antimicrobial property of the gel, secondary infections after bites also heal up rapidly.

Example 16

25 Prophylacticum against skin infections

The treatment of banal scraping wounds should be mentioned as an example of using the gel as a prophylacticum. Primarily the wound is cleansed with water/soap following classic principles. After that, gel  
30 is applied, for example, the gel mentioned in example 1. During the first 24 hours the gel is applied 3 times. After that, the gel is applied once a day until the wounds start to heal. Due to the properties of the dried gel, acting as an elastic, fixed plaster on the wound, a good  
35 protection of the wounds is achieved in between the applications.

Example 17Treatment of Chickenpox (Variola)

5 The gel of example 4 is applied directly onto the individual eruptions on the skin of a patient with chickenpox as soon as possible after the eruption and every 2 to 4 hours thereafter. This immediately soothes the skin, reduces itching, prevents blister formation and renders the patient non-contagious in a shorter period of time. The gels of examples 3 and 10 have similar activity.

10

Example 18Bee Sting and Other Insect Bite Treatment

15 The gel of example 4 is applied direct to the skin at the site of a bee sting as soon as possible after occurrence of the sting. The discomfort and swelling were immediately reduced and itching does not develop. This activity is thought to be due to an effect of the concentrated alcohol in coagulating protein in the insect venom injected into the skin. The gels of examples 3 and 20 10 have similar performance.

CLAIMS

1. A gel form pharmaceutical composition for the treatment of skin disorders comprising a liquid and a high  
5 molecular weight polymer gelling agent dissolved in the liquid, characterised in that the composition comprises more than 90% by weight of a lower alkanol based on the total weight of the composition and less than 10% by weight water based on the total weight of the composition.
- 10 2. A composition according to claim 1 in which the concentration of water in the composition is less than the equilibrium content at temperatures in the range 20-24°C and 50 to 100% relative humidity.
3. A composition according to claim 1 or claim 2  
15 consisting essentially only of the gelling agent, alkanol and water.
4. A composition according to claim 1 or claim 2 comprising further an effective amount of an enhancing agent which enhances the effect of the alcohol in the  
20 treatment of said skin disorder, and being substantially free of other pharmaceutically active components.
5. A composition according to claim 4 which comprises a base, preferably in an amount to give a pH in the range 6 to 9.5, such as an inorganic alkali or an organic base,  
25 preferably selected from sodium and potassium hydroxide and triethylamine.
6. A composition according to any preceding claim in which the alkanol is selected from ethanol, isopropanol and mixtures thereof, and is preferably ethanol.
- 30 7. A composition according to any preceding claim in which the gelling agent is a derivative of cellulose, preferably a cellulose ether, more preferably ethyl hydroxyethyl cellulose.
8. A composition according to any preceding claim in  
35 which the concentration of water is less than 5%, preferably less than 3.3% based on the weight of alkanol plus water.



9. A composition according to any preceding claim contained in a moisture- and moisture vapour-impervious container.
10. Use of a combination of a polymer gelling agent and a lower alkanol in the manufacture of composition containing at least 90% by weight of the lower alkanol and less than 10% water, based on the total composition weight, for treatment of skin infected by a virus.
11. Use according to claim 10 in which the viral infection is of Herpes simplex virus.
12. Use of a combination of a polymer gelling agent and a lower alkanol in the manufacture of a gel form composition containing at least 90% by weight of lower alkanol and less than 5% by weight of water, based on the total composition and gelling agent, for treatment of skin infected by ectoparasites.
13. Use according to any of claims 10 to 12 including the further features defined in any of claims 2 to 9.
14. Use of ethylhydroxyethyl cellulose in the manufacture of a composition for treatment, by topical application, of a skin disorder.
15. A composition containing at least 90% by weight of a lower alkanol and an effective gelling amount of ethylhydroxyethyl cellulose.
16. A composition according to claim 15 for use as a medicament.
17. A method of treatment of infected skin by topical application to the infected area of skin of a gel form pharmaceutical composition comprising a polymer gelling agent, more than 90% by weight of a lower alkanol and less than 10% by weight water based on the total weight of composition.
18. A method according to claim 17 in which the treatment is effective to affect a layer of skin deeper than the stratum corneum.
19. A method according to claims 17 or 18 in which the composition remains in contact with the area affected by

the disorder for a period of at least 2 hours, preferably at least 4 hours, to form a cohesive barrier film of said polymer.

- 5 20. Use of a lower alkanol as a pharmaceutically active ingredient in the manufacture of a composition for use in a method of treatment of disorders of layers of the skin below the stratum corneum by topical application to the skin.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/01025

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 A61K47/10 A61K47/38 A61K9/70

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB,A,1 593 097 (FISONS LIMITED) 15 July 1981 see the whole document ---	1-11, 13, 15-20
X	FR,A,2 613 621 (ROUSSEL-UCLAF) 14 October 1988  see page 3, line 24 - line 36 see page 4, line 24 - line 27 see example 4 ---	1, 2, 6-10, 13-20
X	EP,A,0 320 254 (THAMES PHARMACEUTICAL) 14 June 1989 see claim 1 see page 3, line 31 - page 4, line 3 see page 5, line 20 - line 23 --- -/--	1-9, 13, 16-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

27 July 1995

Date of mailing of the international search report

07.08.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+ 31-70) 340-3016

Authorized officer

Ventura Amat, A

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 95/01025

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,4 671 955 (VICTOR PALINCZAR) 9 June 1987 see claims 1,3,4 see column 3, line 6 - line 12 -----	14

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 95/01025

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 17-19 are directed to a method of treatment of the human body the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/01025

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-1593097	15-07-81	NONE	
FR-A-2613621	14-10-88	AU-B- 616657	07-11-91
		AU-A- 1597688	04-11-88
		DE-A- 3865283	07-11-91
		EP-A, B 0309533	05-04-89
		WO-A- 8807856	20-10-88
		JP-T- 1503461	22-11-89
		QA-A- 8973	30-11-90
		US-A- 4992475	12-02-91
EP-A-0320254	14-06-89	US-A- 5013545	07-05-91
		CA-A- 1334280	07-02-95
		US-A- 5098717	24-03-92
US-A-4671955	09-06-87	AU-B- 591854	14-12-89
		AU-A- 7019287	08-10-87
		DE-A- 3710293	05-11-87
		GB-A, B 2189701	04-11-87
		JP-A- 62265215	18-11-87

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

### **IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**